



# Towards a multiscale crop modelling framework for climate change adaptation assessment


Bin Peng<sup>1,2</sup>✉, Kaiyu Guan<sup>1,2,3,4</sup>✉, Jinyun Tang<sup>5</sup>, Elizabeth A. Ainsworth<sup>4,6,7,8</sup>, Senthod Asseng<sup>9</sup>, Carl J. Bernacchi<sup>3,4,6,7,8</sup>, Mark Cooper<sup>10</sup>, Evan H. Delucia<sup>1,2,3,4,6,8</sup>, Joshua W. Elliott<sup>11</sup>, Frank Ewert<sup>12,13</sup>, Robert F. Grant<sup>14</sup>, David I Gustafson<sup>15</sup>, Graeme L. Hammer<sup>10,16</sup>, Zhenong Jin<sup>17</sup>, James W. Jones<sup>9</sup>, Hyungsuk Kimm<sup>1</sup>, David M. Lawrence<sup>18</sup>, Yan Li<sup>19</sup>, Danica L. Lombardozzi<sup>18</sup>, Amy Marshall-Colon<sup>2,3,4,6,8</sup>, Carlos D. Messina<sup>20</sup>, Donald R. Ort<sup>4,6,8,21</sup>, James C. Schnable<sup>22,23</sup>, C. Eduardo Vallejos<sup>24</sup>, Alex Wu<sup>10,16</sup>, Xinyou Yin<sup>25</sup> and Wang Zhou<sup>1</sup>

**Predicting the consequences of manipulating genotype (G) and agronomic management (M) on agricultural ecosystem performances under future environmental (E) conditions remains a challenge. Crop modelling has the potential to enable society to assess the efficacy of G × M technologies to mitigate and adapt crop production systems to climate change. Despite recent achievements, dedicated research to develop and improve modelling capabilities from gene to global scales is needed to provide guidance on designing G × M adaptation strategies with full consideration of their impacts on both crop productivity and ecosystem sustainability under varying climatic conditions. Opportunities to advance the multiscale crop modelling framework include representing crop genetic traits, interfacing crop models with large-scale models, improving the representation of physiological responses to climate change and management practices, closing data gaps and harnessing multisource data to improve model predictability and enable identification of emergent relationships. A fundamental challenge in multiscale prediction is the balance between process details required to assess the intervention and predictability of the system at the scales feasible to measure the impact. An advanced multiscale crop modelling framework will enable a gene-to-farm design of resilient and sustainable crop production systems under a changing climate at regional-to-global scales.**

Achieving global food security under climate change (CC) is a big challenge facing humanity<sup>1</sup>. Effective adaptation strategies are needed to mitigate the negative CC impacts on crop production and further enhance agricultural resilience. Possible solutions include (1) genetic improvement through plant breeding and genetic engineering of new crops with stress tolerant traits<sup>2,3</sup>, and enhanced resource use efficiencies<sup>4,5</sup>; and (2) adaptive crop management practices<sup>6,7</sup>. As CC impacts on crop production will likely vary substantially in space<sup>2,8</sup>, environment-specific adaptation strategies may confer advantages<sup>6</sup>. Moreover, CC is very likely to shift the target population of environments in which crop cultivars produced by breeding programs will be grown<sup>9</sup> and may limit the rate of genetic gain through breeding programs alone. Considering

both genetic and management improvements can thus expand the opportunities to cope with CC, in which case suitable predictive models that can capture complex interactions among genotype, environment (including climate and edaphic conditions) and management (genetics × environment × management (G × E × M)) in crop development, growth and yield are required<sup>10–12</sup>. Meanwhile, adaptation of crop production to CC should be treated along with environmental sustainability, as the environmental footprint of agriculture may change with adoption of different adaptation strategies over extended areas<sup>13</sup>. Potential genetic and/or management adaptation strategies need to be rigorously assessed for their impacts on both crop production and environmental sustainability in a changing climate (Fig. 1).

<sup>1</sup>Department of Natural Resources and Environmental Sciences, University of Illinois at Urbana-Champaign, Urbana, IL, USA. <sup>2</sup>National Center for Supercomputing Applications, University of Illinois at Urbana-Champaign, Urbana, IL, USA. <sup>3</sup>Institute for Sustainability, Energy, and Environment, University of Illinois at Urbana-Champaign, Urbana, IL, USA. <sup>4</sup>Center for Advanced Bioenergy and Bioproducts Innovation, University of Illinois at Urbana-Champaign, Urbana, IL, USA. <sup>5</sup>Climate Sciences Department, Climate and Ecosystem Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA. <sup>6</sup>Department of Plant Biology, University of Illinois at Urbana-Champaign, Urbana, IL, USA. <sup>7</sup>USDA ARS Global Change and Photosynthesis Research Unit, Urbana, IL, USA. <sup>8</sup>Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, IL, USA. <sup>9</sup>Agricultural and Biological Engineering Department, University of Florida, Gainesville, FL, USA. <sup>10</sup>Centre for Crop Science, Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, Brisbane, Queensland, Australia. <sup>11</sup>Department of Computer Science, University of Chicago, Chicago, IL, USA. <sup>12</sup>Crop Science Group, INRES, University of Bonn, Bonn, Germany. <sup>13</sup>Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, Germany. <sup>14</sup>Department of Renewable Resources, University of Alberta, Edmonton, Alberta, Canada. <sup>15</sup>Independent scientist, St. Louis, MO, USA. <sup>16</sup>Australian Research Council Centre of Excellence for Translational Photosynthesis, The University of Queensland, Brisbane, Queensland, Australia. <sup>17</sup>Department of Bioproducts and Biosystems Engineering, University of Minnesota-Twin Cities, St. Paul, MN, USA. <sup>18</sup>National Center for Atmospheric Research, Boulder, CO, USA. <sup>19</sup>State Key Laboratory of Earth Surface Processes and Resources Ecology, Faculty of Geographical Science, Beijing Normal University, Beijing, China. <sup>20</sup>Corteva AgriScience, Johnston, IA, USA. <sup>21</sup>Department of Crop Science, University of Illinois at Urbana-Champaign, Urbana, IL, USA. <sup>22</sup>Department of Agronomy & Horticulture, University of Nebraska-Lincoln, Lincoln, NE, USA. <sup>23</sup>Center for Plant Science Innovation, University of Nebraska-Lincoln, Lincoln, NE, USA. <sup>24</sup>Horticultural Sciences Department, University of Florida, Gainesville, FL, USA. <sup>25</sup>Centre for Crop Systems Analysis, Department of Plant Sciences, Wageningen University & Research, Wageningen, The Netherlands. ✉e-mail: [binpeng@illinois.edu](mailto:binpeng@illinois.edu); [kaiyug@illinois.edu](mailto:kaiyug@illinois.edu)



**Fig. 1 | Crop modelling plays a central role in assessing agricultural CC adaptation for food security and environmental sustainability.** When optimizing existing and designing new agricultural CC adaptation strategies, lessons from both crop productivity and environmental sustainability assessments should be considered. G, genetics; E, environment; M, management.


### The need for a multiscale crop modelling framework

Crop modelling can help assess the efficacy of agricultural adaptation strategies to CC (Fig. 1). Complex  $G \times E \times M$  interactions<sup>10–12</sup> affecting crop development, growth and yield, and their nonlinearities present a formidable challenge to designing empirical experiments that adequately sample the  $G \times E \times M$  space. Process-based crop models (CMs) offer a unique way to assess the pros and cons of different adaptation strategies because they integrate physical and biological principles to mechanistically simulate the use and allocation of captured resources<sup>14</sup>. In this way, CMs can help optimize existing strategies and assist the design of new strategies under different environmental scenarios (Fig. 1). For management adaptation, many existing studies used CMs to assess the impacts of different practices on crop production<sup>15,16</sup>. However, their potential impacts on environmental sustainability and feedback to the climate remain under-studied, mainly because current CMs lack the required interfaces with land surface, climate and economic models. Crop modelling is an ideal tool to assess the benefits of various genetic solutions in crop improvement<sup>17</sup>. However, assessments of potential genetic solutions by most existing CMs are restricted due to their limited capabilities to simulate genetic variation and biochemistry<sup>18,19</sup>. Some recent model developments have uncovered opportunities to develop an integrated modelling framework to address questions about genetic improvement under CC. For instance, CMs were improved by considering the variation in relevant adaptive traits to drought in maize<sup>20</sup> as well as the connection between CMs and genomic prediction<sup>21,22</sup>. Although geospatial simulations of the impact of crop genetic improvements on yield exist<sup>23</sup>, assessments of their impact on environmental sustainability under

current and future climates are still missing. To meet these demands for assessing CC adaptation strategies for both crop productivity and environmental sustainability, and assisting in designing new strategies, a multiscale (from gene to globe) crop modelling framework should be developed using systems approaches. Such a framework can explicitly integrate small-scale mechanisms with multi-sectoral impacts of different adaptation strategies at larger scales (Fig. 2; Table 1).

### Going to gene scale

Incorporating the principles of genetics and genomics into CMs will enable genotype-to-phenotype prediction to advance crop breeding and genetic engineering<sup>18,24,25</sup>. Two approaches, referred to as ‘top-down’ and ‘bottom-up’ approaches, can be employed to leverage our collective understanding of genetic controls of physiological processes to enhance phenotypic prediction<sup>26–28</sup>. The top-down approach<sup>25,29</sup> follows the philosophy of ‘modelling plant hormone action without modelling the hormones’<sup>30</sup>, and strives to capture system dynamics and phenotypic consequence from genetic variation at the whole-plant scale, with a relatively coarse granularity but robust prediction accuracy<sup>25</sup>. The simplification made in the top-down approach balances phenotyping capabilities<sup>31</sup>, propagation of measurement errors and accuracy, which are required to inform a decision. To effectively connect complex traits (such as yield) with genetic regulation, CMs should include detailed physiological processes<sup>10,29,31,32</sup>. Some cultivar-specific model parameters in CMs are predicted by different mathematical approaches, from simple statistical models based on multiple loci<sup>32–35</sup> to whole genome association prediction using Bayesian and machine learning approaches<sup>21,22,36</sup>. Successful examples of this top-down



**Fig. 2 | A conceptual illustration of the multiscale crop modelling framework. a,** Opportunities to advance the multiscale crop modelling. **b,** Models operating at different scales involved in the multiscale crop modeling framework. **c,** Data collection to support model development and evaluation at varied scales.

approach include reasonable predictions with quantitative trait loci models of maize leaf elongation response to soil water deficit and high evaporative demand<sup>33,34</sup>, and the transition to reproductive growth in several other crops<sup>32,37</sup>. CMs that incorporated functional relationships between environmental variables and traits for the genetic variation in yield and agronomic performance<sup>22</sup> have been successfully used in breeding drought-tolerant maize hybrids for the US Corn Belt<sup>3</sup>.

The bottom-up approach uses systems biology to integrate plant biological processes operating at different temporal and spatial scales<sup>26,38,39</sup>. For example, the *e*-photosynthesis model<sup>40</sup>, which simulates individual photosynthesis-related metabolic reactions and their major regulatory mechanisms, is a typical bottom-up model that may potentially link high throughput genomic data with observable macroscopic phenotypes. By explicitly simulating gene network regulation, metabolic reactions and metabolite transport, the bottom-up models, once integrated to the whole-plant scale, could enable testing the impact of genetic manipulation on different proteins (such as photosynthetic enzymes) and identify targets for crop genetic improvements<sup>24</sup>. However, the bottom-up approach is limited to single component processes (for example, photosynthesis)<sup>40,41</sup> or model plants (for example, *Arabidopsis*) that have relatively lower complexity and have been better characterized than conventional crop species<sup>42</sup>. Mechanistic models that include all component processes controlling the forms and functions of crops are being investigated<sup>43</sup>. Although the bottom-up approach can provide a more direct connection with the underpinning of genetic

architecture, several key challenges need to be met before scaling up to the whole-plant level. These include how to integrate the component modules and whether the integrated models can give robust phenotypic responses to genetic changes under varied environmental and management conditions, which may be challenging in applications that require a higher prediction accuracy due to error propagation and inability to properly phenotype and initialize such models.

Given the different properties of top-down and bottom-up approaches, integrating these two approaches may provide an opportunity to merge their advantages. Such integration will require transdisciplinary cooperation to harness collective knowledge in the form of functional models that capture essential features of the underlying biology while retaining the required predictability. Comparing these two approaches for genotype-to-phenotype prediction under varied environmental conditions will help identify potential model deficiencies and corresponding opportunities for improvement. To make them operationally feasible, the fine-grained bottom-up models must be simplified in mechanistic representations while maintaining predictive capability of system dynamics for large-scale applications<sup>44</sup>. One possibility is to build surrogate models of the bottom-up models which can statistically emulate the behaviours of the latter but with greater computational efficiency. This is critical for some applications requiring many model runs, such as high-resolution simulation over large areas and the Monte Carlo-based parameter estimation. Surrogate models for selected processes could also be combined with top-down models to build a hybrid model to simulate the whole-plant-level impact of crop

**Table 1 | A summary of recommended actions to advance multiscale crop modelling**

Directions or opportunities	Recommended actions	Process understanding	Model development and evaluation	Data collection and model-data integration
Going to gene scale	Comparing top-down and bottom-up approaches for genotype-to-phenotype simulation	✓	✓	✓
	Integrating top-down and bottom-up approaches to represent genetic traits in CMs		✓	✓
Going to global scale	Interfacing CMs with large-scale land surface, climate and economic models	✓	✓	✓
	Scaling the surface heterogeneity from field to regional or global scale		✓	
Better representation of physiological responses to CC	Simulating coupled soil-root-canopy-atmosphere water transfer driven by energy balances	✓	✓	
	Improving the stomatal and intra-leaf diffusional conductance models	✓	✓	
	Improving the simulation of responses of carbon or nitrogen source-sink relationship to stresses	✓	✓	
	Developing mechanistic models for ozone stress	✓	✓	
	Simulating the root growth and metabolism under oxygen deficiency	✓	✓	
Better representation of crop management practices	Simulating coupled carbon-nitrogen-phosphorus cycles in CMs	✓	✓	✓
	Simulating microorganism-root interactions in CMs	✓	✓	✓
	Representing more management practices in large-scale CMs	✓	✓	✓
	Simulating stresses from crop pests and diseases as well as weed competition on crop growth	✓	✓	✓
	Improving simulation of fate and transport of pesticide across landscapes		✓	✓
Closing the data gaps	Collecting more site-level experimental data following standardized protocols			✓
	Conducting multi-dose experiments for observed crop responses to CC factors			✓
	Collecting more soil profile data to improve the gridded soil products			✓
	Enriching management data through working with farmers and government agencies, and using crowdsourcing and remote sensing technologies			✓
Model-data integration	Evaluating CMs using eddy-covariance flux data		✓	✓
	Evaluating CMs in simulating the emergent relationships inferred from data		✓	✓
	Spatially explicit calibrating CMs using remote sensing data as constraints		✓	✓

trait manipulations at finer (gene and metabolic) scales effectively and efficiently. Meanwhile, lessons from the top-down approach may help component integrations in the bottom-up approach, as top-down models provide robust physiological overviews of crop growth and development processes. The top-down approach can also be used in designing better structures for those simplified mechanistic models or surrogate models. Such integration provides a promising approach to represent genetic traits in CMs for breeding and genetic engineering research—a research domain which is still in its infancy<sup>24</sup>.

### Going to global scale

Reliable large-scale simulations to assess agricultural CC adaptation are needed because decisions about adaptation are usually made

at a broader scale, beyond that which the traditional CMs were developed and tested (that is, plot-to-field scale)<sup>45</sup>. Current CMs are mainly designed for yield simulation at the field scale, excluding or implicitly considering processes related to surface fluxes of water, energy, carbon and nutrients. For these reasons, current CMs appear unsuitable for evaluating the environmental impacts of agricultural CC adaptation at large scales. Meanwhile, land surface models (LSMs) and their counterparts in Earth system models (ESMs) focus more on surface and subsurface hydrology, surface energy balance and biogeochemical cycling processes<sup>46</sup>. Many LSMs can be coupled to an atmospheric model, a river model, an ocean model (via the river model) and, occasionally, even to an integrated assessment model. These couplings could enable both assessment of the broader role agriculture plays in the Earth system and

studies regarding food security, and environmental quality under socioeconomic development, technological innovations and CC. Although some LSMs have incorporated crop growth and a number of plant physiological mechanisms, such as stomatal and biochemical limitations to photosynthesis and plant hydraulics, crop modelling schemes in ESMs require development of crop-specific processes that are better resolved by field-scale agronomic CMs<sup>17</sup>. Most ESM-based CMs do not incorporate sufficient details describing crop phenology, organ development and their responses to environmental stresses; therefore, they cannot always capture observed variation in crop yields. These inter-model differences occur because field-scale agronomic CMs and LSMs, or ESMs, were originally developed for different goals. LSMs or ESMs can become powerful tools to simulate the ‘food–energy–water’ nexus<sup>48</sup>, the crux of both food security and environmental sustainability, if essential biological functionalities related to both surface energy–water–carbon fluxes and crop growth and yield are included.

Regional-to-global CC adaptation assessment requires a systems approach to understand interactions among different elements of the climate–agriculture–environment system, which can be leveraged to increase overall system efficiency, resilience and sustainability. One possible research pathway is to integrate field-scale crop modelling capabilities with LSMs or ESMs<sup>47,49,50–52</sup> making CMs a sub-module of larger system models. This integration will enable both CMs and the host LSMs to complement and benefit from each other. CMs can take advantage of the more advanced biophysical and biogeochemical representations in LSMs to help improve the simulation of crop growth and yield at regional-to-global scales. Similarly, LSMs can benefit from the explicit and mechanistic representation of crop growth to improve surface energy–water–carbon flux simulation over agricultural land. For regional-to-global scale applications, trade-offs between model complexity and computational efficiency need to be considered<sup>29</sup>. For example, when incorporating genetic models, the coarse-grained top-down approach<sup>33</sup> may be preferred for its better computational efficiency and reliability compared to the fine-grained bottom-up approach. In addition, the three-dimensional representation of leaf anatomy<sup>44</sup> as well as root and canopy architectures<sup>53,54</sup> in structural–functional models<sup>55</sup> may also need to be simplified in LSMs for regional-to-global scale applications<sup>44,56</sup>.

One challenge when integrating CMs with large-scale models is scaling the heterogeneity over cropland<sup>45,57</sup>. Most large-scale crop simulations still use field-scale CMs with spatially aggregated input data (soil, weather and management), such as those in the Global Gridded Crop Model Intercomparison<sup>58,59</sup> of the Agricultural Model Intercomparison and Improvement Project (AgMIP; <https://agmip.org/>)<sup>60</sup>. However, such aggregation may lead to biases in both simulated yield and its variability depending on regional characteristics, CMs and spatial resolution of simulation<sup>45,61</sup>. Finer resolutions could somewhat mitigate the problem but would dramatically increase the computational burden, especially when using ESM-based CMs at regional-to-global scales. Recent advances in representing land surface heterogeneity in ESMs also shed light on the scaling issue in crop modelling. As more data on field-scale elevation, soil, yield and management become available, high-resolution cropland grids (~30 m) can be clustered into several tiles based on the multi-dimensional characteristic space<sup>62</sup>. Simulations over the clustered tiles within one coarse grid can then be mapped back to high-resolution grids through post-processing. Necessary adjustments to model structures may be needed when adding this scaling complexity into CMs, especially for those using a single soil column for all cropland within one coarse grid.


Integrating CMs with large-scale models enables a consistent and systematic assessment of the adaptation impacts on multiple sectors of the Earth system at regional-to-global scales. Possible applications include determining how genetic modifications of crop

growth, vigour, stress response or resource use efficiency influence water–energy–carbon cycles at regional-to-global scales and their potential feedback to the climate. These large-scale impacts of possible genetic improvements can be assessed using ESMs with embedded CMs to guide crop breeding, designing sustainable cropping systems and their placement on the landscape. Likewise, adaptive changes in the amount and timing of irrigation<sup>7</sup> as well as nitrogen fertilization<sup>63</sup> in response to CC may have economic and environmental implications that can be assessed with the coupled crop, economic and environmental models<sup>64</sup>.

### Better representation of physiological responses to CC

Realistic physiological responses to CC in CMs are essential for rigorous CC adaptation assessment<sup>65</sup>. Multi-model intercomparisons, like AgMIP<sup>60</sup>, show that current CMs can simulate comparable yields under current climate and management conditions<sup>66–69</sup>. However, those models diverge significantly in predicting future yield under common CC scenarios<sup>67,70,71</sup>, mainly due to their diverging assumptions about the underlying physiology of how crop growth will respond to changes in various environmental factors, including carbon dioxide concentration ( $[CO_2]$ )<sup>67,72,73</sup>, temperature<sup>67,73,74</sup>, soil water and nutrient availability<sup>68</sup>. The divergence issue becomes more complicated with their significant differences in representing combined effects of different environmental stresses on plant development and biomass production<sup>31,66</sup>. Multi-model ensemble can lead to more robust model predictions with a potentially narrower uncertainty range than an individual model projection<sup>67–69,71,75,76</sup>. However, since most CMs have limited capability to simulate the combined effects of multiple CC factors on crop growth, confidence in such multi-model projections is reduced, especially when considering climate extremes (droughts, heatwaves and flooding) and their interactions.

Higher  $[CO_2]$  alone is expected to benefit crop production. Free air  $CO_2$  enrichment (FACE) experiments show that elevated  $[CO_2]$  may increase net carbon assimilation and plant growth for  $C_3$  crops and for  $C_4$  crops under water-limited conditions<sup>77,78</sup>. Elevated  $[CO_2]$  also reduces stomatal conductance and potentially enhances water use efficiency<sup>79–81</sup>. However, this does not necessarily lower the drought risk, since elevated  $[CO_2]$  may stimulate leaf growth and hence lead to more water consumption in some crop species, such as soybean<sup>82,83</sup>. Moreover, reduced canopy cooling associated with lower transpiration rates under elevated  $[CO_2]$  could also increase canopy temperature<sup>80</sup>. The magnitudes of these  $CO_2$  effects vary depending on crop species and cultivar<sup>84,85</sup> as well as environmental conditions because the  $CO_2$  effect interacts with temperature, water, nitrogen and ozone stress conditions<sup>79,86–88</sup>. The modelling community has attempted to reconcile model predictions with these observed results, but inconsistencies between model simulation and observations remain<sup>72</sup>. Part of this model–data discrepancy may come from different approaches CMs use to represent plant physiological responses to elevated  $[CO_2]$ . For example, CMs may simulate canopy-level assimilation using either radiation use efficiency (RUE)<sup>89</sup> or photosynthetic biochemical limitation approaches<sup>17,90</sup>. In the RUE approach, the  $CO_2$  fertilization effect on assimilation is simulated through a multiplicative modifier to RUE, which is often empirically derived from limited experimental observations<sup>91</sup>. A better understanding of the biochemistry-related responses of leaf-scale stomatal and intra-leaf diffusional conductance, autotrophic respiration to elevated  $[CO_2]$  and their scaling to canopy scale is needed for improved model performance<sup>86,92</sup>. Further, long-term effects of elevated  $[CO_2]$  on crop growth differ from short-term effects due to photosynthetic acclimation, and current CMs differ in their abilities to translate short-term responses to long-term effects<sup>93</sup>. Coupling  $CO_2$  response to nutrient limitation through product inhibition and understanding the source–sink relationship in carbon and nutrient allocation under varied environmental



**Fig. 3 | Temperature free-air controlled enhancement experiment for soybean in Illinois, USA.** **a**, Experiment design for the multi-level temperature treatment (+1.5 °C, +3.0 °C, +4.5 °C and +6.0 °C).  $T_c$ , canopy temperature. **b**, Different physiological mechanisms depicting the responses of soybean growth and yield to higher temperature. AGB, above-ground biomass; HI, harvest index. **c**, A conceptual temperature response function from the field measurement that can be used to constrain CMs, enabling models to reproduce the right yield response with the right mechanism.

conditions are essential for CMs to capture the yield responses to elevated  $[\text{CO}_2]$ <sup>94</sup>.

Chronic or acute exposure of crops to tropospheric ozone can decrease photosynthesis, alter carbon allocation and reduce yield quantity and quality<sup>95</sup>. As ozone damage occurs at the cellular level, CMs need to explicitly represent leaf photosynthesis biochemistry, stomatal conductance and daily or sub-daily canopy growth and biomass partitioning to simulate ozone impacts on crop growth, yield and its interaction with other environmental factors. However, current CMs seldom consider these physiological mechanisms of ozone stress on crop growth<sup>96–98</sup>, and ozone-induced yield impact assessments at regional-to-global scales mainly rely on empirical or semi-empirical modelling approaches<sup>95</sup>. Future studies should prioritize the development and benchmark of a mechanistic algorithm for ozone stress on crop growth using high-quality data from ozone filtration and fumigation experiments conducted in open top chambers or FACE sites<sup>95,99,100</sup>.

Since extreme events, such as heatwaves, droughts and extreme rainfall events, are projected to increase under future climate scenarios<sup>101</sup>, representing their impact on crop growth and productivity in CMs is essential for projecting future food security<sup>102</sup>. Heat stress affects grain setting during anthesis and grain size during the grain filling period<sup>103–105</sup>, and therefore can directly lead to yield reduction. Asymmetric impacts of high day and night temperatures have been observed for some crop species, particularly rice<sup>106,107</sup>. Drought stress can reduce stomatal conductance, downregulate photosynthesis, slow plant development and growth, cause earlier leaf senescence and reduce yields<sup>103</sup>. The impacts of heat and drought stresses are growth-stage-dependent. For example, greater yield loss is usually observed if crops are stressed from heat and drought during meiosis, anthesis and grain filling stages<sup>108</sup>. Moreover, heat and drought stresses usually occur concurrently and interact with each other, increasing the overall stress for crop growth<sup>109,110</sup>. The crop breeding community has been targeting new cultivars that tolerate both drought and heat stresses over the last two decades. However, current CMs show suboptimal performance in simulating the impacts of drought-heat interacting stresses<sup>111,112</sup>, mainly due to complexity in simulating plant physiological responses to such complex abiotic conditions. Deficiencies in model structure can also lead to uncertainty in assessing heat stress impacts on crop yield and understanding the relative importance of drought as compared to heat stress<sup>113,114</sup>. Examples of this problem include using air temperature instead of canopy temperature to quantify heat stress<sup>115,116</sup>, and

lack of canopy energy budget closure and plant hydraulics in most traditional CMs. Compared with drought and heat stress, negative impacts of excess rainfall or waterlogging on crop growth have received even less attention in current CMs<sup>117,118</sup>, although it may lead to comparable yield losses<sup>119</sup>. Root growth and metabolism under oxygen deficiency should be explicitly simulated in CMs to quantify the oxygen limitations on active nutrient uptake and crop growth. Understanding and implementing these physiological responses to changing environmental factors in CMs should be prioritized and would benefit model-assisted design of CC adaptation strategies.

### Better representation of crop management practices

Crop management practices are the actions farmers take to ensure crop productivity. Current field-scale CMs generally have specific crop management modules which can simulate parts of the impacts from some management practices. However, opportunities for improvement are abundant, especially considering these CMs simulate largely diverged sensitivities of crop yield, soil carbon and nitrogen fluxes to management practices<sup>120,121</sup>. An emerging priority is to improve the model representation of management impacts on soil biogeochemical cycling. Unlike natural ecosystems, biogeochemical cycling in farmland soils is directly affected by crop management practices, and it is a key determinant of crop productivity and CC mitigation potential of agroecosystems. Improved representation of plant–soil–microbial interactions and the impacts of management practices on such interactions could improve simulating coupled carbon–nitrogen–phosphorus cycles in cultivated soil<sup>121</sup>. Recent advancements in soil biogeochemistry and ecosystem modelling could be combined and leveraged<sup>122–125</sup>. Additionally, most previous model-ensemble-based assessments of CC impacts on crop production reinitialize soil water and nutrient conditions in each growing season to solely focus on impacts from climate variability<sup>66,67,126,127</sup>. Only recently have continuous multi-year simulations been used to assess the legacy impacts of management practices on both crop yield and environmental sustainability (such as soil health and greenhouse gas emissions)<sup>128–131</sup>. Inclusion of regionally representative crop management remains a priority for ESM-based CMs<sup>132</sup>. Current ESMs consider a limited number of crop management practices, such as irrigation and nitrogen fertilization<sup>49</sup>. Other crop management practices (such as tillage, crop rotation and cover crops) still need to be implemented. A challenge to implement all of them represents the large data gaps in spatiotemporal characterization of these crop management practices.

Improving pest and disease management simulation in CMs is another priority as crop yield loss from these processes is projected to increase with CC<sup>133–135</sup>. Few CMs simulate the stress from pests and diseases, which, if not well controlled, can be a key determinant of yield gaps. Process-based simulation of crop pest and disease management requires at least three sub-modules: population dynamics, injury and damage, and management action. Population dynamics models have been well developed in the epidemiological community. Linking those population models with crop growth models is the first step to address the complex interactions between crop and pest or pathogen dynamics<sup>14,136</sup>. However, this coupling requires a better understanding of the injuries and damage mechanisms. For example, modelling corn borer injury to xylem vessels, and hence water uptake, requires a hydraulic-driven model of soil–root–canopy–atmosphere water transfer along potential gradients through a series of hydraulic resistances. Pest and disease control actions, such as pesticide use, should be simulated by explicitly considering their impacts on population dynamics and metabolism rates of pests and diseases. Improving the simulation of fate and transport of pesticides in the soil–crop–atmosphere continuum as well as in rivers would benefit environmental sustainability assessment<sup>137,138</sup>. With these advancements, potential benefits in both crop production and environmental sustainability of using genetic mechanisms for protection against pest and disease damage could also be assessed using the multiscale modelling framework.

### Closing the data gaps

Observations from individual sites or site networks have been widely used to evaluate and improve field-scale CMs<sup>139</sup>. However, current field experiments have limited coverage of growth conditions (climate, soil and crop management practices). More site-level data collection following standardized protocols for model evaluation and improvement is needed<sup>140</sup>. In addition, current field data generally have limited treatment levels for [CO<sub>2</sub>], temperature or [O<sub>3</sub>]. Many previous studies have evaluated CMs mainly for their accuracy in simulating target variables such as crop yield, above-ground biomass and leaf area index. Any future projections on crop production should consider that simulating the emergent relationships is as important as, or perhaps more important than, simulating the ‘right magnitude’ of any particular target variable in CMs<sup>71</sup>. Examples of such emergent relationships are the nonlinear yield response to temperature<sup>141</sup> and kernel number response to plant growth<sup>20</sup>. Focusing on functional relationships could minimize or prevent CMs from predicting right values due to the wrong mechanism<sup>65</sup>.

Progress in science and technology is an iterative process whereby experimentation will inform modelling and vice versa. The effectiveness of integrating field research programs with model development and evaluation efforts at the process level across environmental gradients could be increased by designing standardized protocols to collect field data. Future field experiments should be designed to facilitate model development and validation of the ‘responses’ of CMs to various environmental conditions and stressors. A good example is the multi-level temperature free-air controlled enhancement experiment in Illinois (USA), which collected data for response curves of soybean growth over multiple elevated temperatures (+1.5 °C, +3.0 °C, +4.5 °C and +6.0 °C), instead of a single-dose response value (Fig. 3). Both biometric and leaf gas exchange measurements were collected periodically across multiple growing seasons, providing valuable data for mechanism understanding and model testing.

Data gaps are even larger for crop modelling at regional-to-global scales, especially on soil and crop management characteristics. One major uncertainty is in soil characterization<sup>142</sup>. Though global soil databases exist, their collected soil profiles are insufficient to account for the large heterogeneity in soil characteristics. Moreover,

the spatial distribution of soil profiles is largely biased. Traditional soil maps are vector or polygon products linking dominant soil types for different soil horizons in the same soil column as tabular characteristic data without any documented uncertainty. The variability within each soil map unit polygon is largely neglected depending on the map scales<sup>142</sup>. Regional-to-global gridded soil products have recently become available from spatial interpolation with other environmental covariates<sup>143,144</sup>, but generalization of the soil–environment prediction models is still poor. Besides, soil parameters from these gridded products are not enough to drive CMs in many regions<sup>145</sup>. As an example, the organic C:N:P ratio is a key attribute currently lacking in those soil products. For large-scale crop modelling, the dominant soil-type approach is still widely used to derive soil parameters, which inevitably introduces uncertainties when either upscaling soil types or applying pedotransfer functions at larger scales<sup>146</sup>, sometimes greater than uncertainties from climate aggregation<sup>61</sup>. Moreover, soil characteristics over cultivated areas are actually not static but responsive to CC, land use and management practices<sup>129</sup>. Ignoring this variability may lead to inconsistency in existing soil maps as they are produced from soil surveys conducted at different times. Advancements in new soil sensor development, in situ soil data collection methods and fast soil testing techniques would help collect denser soil datasets with greater frequencies, which will ultimately improve the quality of gridded soil data needed for crop modelling at large scales.

Insufficient agricultural management data at regional-to-global scales also hinders spatiotemporally explicit representation of agricultural management practices in CMs<sup>147</sup>. For example, current global simulation efforts usually use one single reference cultivar or several region-specific cultivars for each crop type without explicit quantification of uncertainties from intra-species variability<sup>58,148</sup>. The spatial distribution and interannual variability in land devoted to specific cultivars of crop species remains poorly quantified. Some of these data gaps are related to strongly held desires for data privacy among farmers. Finding effective ways to work with farmers could be the key to narrow these important data gaps. University extension services and regional farmer groups can play critical roles to facilitate farmer data sharing, research dissemination and farmer benefit from new technologies. Governmental policies on data sharing may also help harness the power of agricultural data collected by public agencies<sup>149</sup>. In addition, crowdsourcing data<sup>150</sup>—for example, crop phenological stages collected by voluntary observer network<sup>151</sup>—can complement such information. Another promising solution is to map agronomic management practices from space. Previous studies have demonstrated the potential of remote sensing in mapping crop species<sup>152</sup>, irrigation area and intensity<sup>153</sup>, and agronomic conservation practices (for example, cover crops and field tillage)<sup>154,155</sup> at large scales. Important crop phenological states, such as sowing, emergence, anthesis, tasseling and silking (for maize)<sup>156–158</sup> can also be derived from remote sensing, which may assist setting management configurations (for example, sowing or harvesting dates and cultivar maturity groups) in CMs.

### Model-data integration

Reconciling model simulations with observations is a natural process to advance model improvements. Besides biomass and yield observations, there is still a largely unrealized opportunity for testing CMs against eddy covariance measurements of water, energy and CO<sub>2</sub> exchange under contrasting environmental conditions. These tests will help identify crop responses to changes in diverse environmental conditions, yet have been largely overlooked by the crop modelling community. A wealth of eddy covariance data has been accumulated at tens of cropland sites and compiled in either regional or global flux networks<sup>159</sup>. It was not until recently that high-frequency eddy covariance observations were used to benchmark simulations of multiple CMs<sup>160</sup>. Besides

conventional model-data comparisons, there is a greater need for evaluation and calibration of the response and emergent relationships in model simulations. Experimental response relationships (such as measured temperature responses from experiments in Fig. 3) can be used to evaluate and constrain CMs at site scale<sup>161</sup>. Simulated yield responses to environmental variability at regional scales can further be compared and calibrated using statistical model-inferred yield responses<sup>141,162</sup>, and, in turn, the insights from statistical models can help improve process-based CMs for better CC adaptation assessment<sup>163,164</sup>. With these ‘responses’ as constraints, CMs could be closer to ‘delivering right simulations with right mechanisms’. The constrained model would enable identifying bottleneck processes for yield production under current and future climate scenarios at regional-to-global scales, and the resultant assessment can guide crop genetic improvement. Remote sensing observations provide another promising opportunity for spatially explicit calibration of CMs at regional-to-global scales. Complementary information about crop growth and stress across different spectral bands<sup>165</sup>, with broad spatial coverage and increasingly finer spatiotemporal resolution of remote sensing data<sup>166</sup>, allow explicit incorporation of satellite information to improve regional crop simulation. Satellite-derived crop phenology stages<sup>156–158</sup>, leaf area index<sup>167</sup>, water use<sup>168</sup> and yield<sup>169</sup> can be used as model constraints<sup>170</sup>.

### Concluding vision

The research community is actively seeking CC adaptation strategies that can ensure future crop production and environmental sustainability. As conducting trial experiments in all environmental conditions is not feasible and some future environmental conditions do not yet exist, crop modelling is a critical tool for hypothesis testing and scenario analysis under varied environmental conditions. Scenario-based model simulations can potentially guide genetic improvement and management adaptation plans. To fulfil this potential, crop modelling must extend beyond its current scales (organ-to-field scales) to both smaller (gene-to-cell) and larger (regional-to-global) scales. Although not all scales must necessarily be included in a single model for all applications, coherent process representations and standardized interfaces to other scales should be developed to facilitate assembling modelling solutions suitable for specific demands. The development of a multiscale crop modelling framework requires multidisciplinary knowledge, including (but not limited to) genetics, genomics, molecular systems biology, plant physiology, agronomy, soil science, agroecology, hydrology, biogeochemistry, climate, Earth system science, remote sensing and engineering<sup>171</sup>. Transdisciplinary collaboration and convergent research approaches are needed to achieve such a multiscale crop modelling paradigm, which will allow for more robust and useful predictions of crop production under CC along with its environmental implications. Broader collaboration among crop modellers across different communities, and promotion of knowledge exchange, data sharing, open-source coding and community-based model development, benchmarking and intercomparison, are fruitful paths forward to success. Several global partnerships, such as AgMIP<sup>60</sup>, Research Program on Climate Change, Agriculture and Food Security (CCAFFS) of the Consultative Group for International Agricultural Research (CGIAR)<sup>172</sup> and Crops *in silico* (Cis)<sup>38,39</sup> serve as guiding examples.

Received: 8 October 2019; Accepted: 24 February 2020;  
Published online: 15 April 2020

### References

- Long, S. P., Ainsworth, E. A., Leakey, A. D., Nösberger, J. & Ort, D. R. Food for thought: lower-than-expected crop yield stimulation with rising CO<sub>2</sub> concentrations. *Science* **312**, 1918–1921 (2006).
- Asseng, S. et al. Climate change impact and adaptation for wheat protein. *Glob. Change Biol.* **25**, 155–173 (2019).
- Cooper, M., Gho, C., Leafgren, R., Tang, T. & Messina, C. Breeding drought-tolerant maize hybrids for the US corn-belt: discovery to product. *J. Exp. Bot.* **65**, 6191–6204 (2014).
- Glowacka, K. et al. Photosystem II Subunit S overexpression increases the efficiency of water use in a field-grown crop. *Nat. Commun.* **9**, 868 (2018).
- Kromdijk, J. et al. Improving photosynthesis and crop productivity by accelerating recovery from photoprotection. *Science* **354**, 857–861 (2016).
- Hammer, G. L. et al. Crop design for specific adaptation in variable dryland production environments. *Crop Pasture Sci.* **65**, 614–626 (2014).
- Zhao, G. et al. The implication of irrigation in climate change impact assessment: a European-wide study. *Glob. Change Biol.* **21**, 4031–4048 (2015).
- Lobell, D. B. et al. Prioritizing climate change adaptation needs for food security in 2030. *Science* **319**, 607–610 (2008).
- Chapman, S. C., Hammer, G. L., Butler, D. G. & Cooper, M. Genotype by environment interactions affecting grain sorghum. III. Temporal sequences and spatial patterns in the target population of environments. *Aust. J. Agr. Res.* **51**, 223–234 (2000).
- Wang, E. et al. Improving process-based crop models to better capture genotype×environment×management interactions. *J. Exp. Bot.* **70**, 2389–2401 (2019).
- Chenu, K. et al. Contribution of crop models to adaptation in wheat. *Trends Plant Sci.* **22**, 472–490 (2017).
- Hammer, G., McLean, G., Doherty, A., van Oosterom, E. & Chapman, S. in *Sorghum: State of the Art and Future Perspectives Agronomy Monographs* Ch. 17 (American Society of Agronomy and Crop Science Society of America, 2016).
- Hunter, M. C., Smith, R. G., Schipanski, M. E., Atwood, L. W. & Mortensen, D. A. Agriculture in 2050: recalibrating targets for sustainable intensification. *BioScience* **67**, 386–391 (2017).
- Jones, J. W. et al. Toward a new generation of agricultural system data, models, and knowledge products: State of agricultural systems science. *Agr. Syst.* **155**, 269–288 (2017).
- Challinor, A. J., Ewert, F., Arnold, S., Simelton, E. & Fraser, E. Crops and climate change: progress, trends and challenges in simulating impacts and informing adaptation. *J. Exp. Bot.* **60**, 2775–2789 (2009).
- Hernandez-Ochoa, I. M. et al. Adapting irrigated and rainfed wheat to climate change in semi-arid environments: management, breeding options and land use change. *Eur. J. Agron.* **109**, 125915 (2019).
- Wu, A., Hammer, G. L., Doherty, A., von Caemmerer, S. & Farquhar, G. D. Quantifying impacts of enhancing photosynthesis on crop yield. *Nat. Plants* **5**, 380–388 (2019).
- Yin, X., van der Linden, C. G. & Struik, P. C. Bringing genetics and biochemistry to crop modelling, and vice versa. *Eur. J. Agron.* **100**, 132–140 (2018).
- Rötter, R. P., Tao, F., Höhn, J. G. & Palosuo, T. Use of crop simulation modelling to aid ideotype design of future cereal cultivars. *J. Exp. Bot.* **66**, 3463–3476 (2015).
- Messina, C. D. et al. On the dynamic determinants of reproductive failure under drought in maize. *in silico*. *Plants* **1**, diz003 (2019).
- Messina, C. D. et al. Leveraging biological insight and environmental variation to improve phenotypic prediction: integrating crop growth models (CGM) with whole genome prediction (WGP). *Eur. J. Agron.* **100**, 151–162 (2018).
- Cooper, M., Technow, F., Messina, C., Gho, C. & Totir, L. R. Use of crop growth models with whole-genome prediction: application to a maize multi-environment trial. *Crop Sci.* **56**, 2141–2156 (2016).
- Sinclair, T. R., Soltani, A., Marrou, H., Ghanem, M. & Vadez, V. Geospatial assessment for crop physiological and management improvements with examples using the simple simulation model. *Crop Sci.* **59**, 1–9 (2019).
- Chang, T.-G., Chang, S., Song, Q.-F., Perveen, S. & Zhu, X.-G. Systems models, phenomics and genomics: three pillars for developing high-yielding photosynthetically efficient crops. *in silico* *Plants* **1**, diy003 (2019).
- Hammer, G. et al. Models for navigating biological complexity in breeding improved crop plants. *Trends Plant Sci.* **11**, 587–593 (2006).
- Minorsky, P. V. Achieving the *in silico* plant. Systems biology and the future of plant biological research. *Plant Physiol.* **13**, 404–409 (2003).
- Hammer, G. L., Sinclair, T. R., Chapman, S. C. & van Oosterom, E. On systems thinking, systems biology, and the *in silico* plant. *Plant Physiol.* **134**, 909–911 (2004).
- Yin, X. & Struik, P. C. Modelling the crop: from system dynamics to systems biology. *J. Exp. Bot.* **61**, 2171–2183 (2010).
- Hammer, G. L. et al. Adapting APSIM to model the physiology and genetics of complex adaptive traits in field crops. *J. Exp. Bot.* **61**, 2185–2202 (2010).
- de Wit, C. T. & Penning de Vries, F. W. T. Crop growth models without hormones. *Neth. J. Agr. Sci.* **31**, 313–323 (1983).
- Parent, B. & Tardieu, F. Can current crop models be used in the phenotyping era for predicting the genetic variability of yield of plants

- subjected to drought or high temperature? *J. Exp. Bot.* **65**, 6179–6189 (2014).
32. Messina, C. D., Jones, J. W., Boote, K. J. & Vallejos, C. E. A gene-based model to simulate soybean development and yield responses to environment Florida agricultural experiment station, journal series no. R-11017. *Crop Sci.* **46**, 456–466 (2006).
  33. Chenu, K. et al. Simulating the yield impacts of organ-level quantitative trait loci associated with drought response in maize: a ‘gene-to-phenotype’ modeling approach. *Genetics* **183**, 1507 (2009).
  34. Reymond, M., Muller, B., Leonardi, A., Charcosset, A. & Tardieu, F. Combining quantitative trait loci analysis and an ecophysiological model to analyze the genetic variability of the responses of maize leaf growth to temperature and water deficit. *Plant Physiol.* **131**, 664 (2003).
  35. Gu, J., Yin, X., Zhang, C., Wang, H. & Struijk, P. C. Linking ecophysiological modelling with quantitative genetics to support marker-assisted crop design for improved yields of rice (*Oryza sativa*) under drought stress. *Annal. Bot.* **114**, 499–511 (2014).
  36. Kadam, N. N., Krishna Jagadish, S., Struijk, P. C., der Linden, C. & Yin, X. Incorporating genome-wide association into eco-physiological simulation to identify markers for improving rice yields. *J. Exp. Bot.* **70**, 2575–2586 (2019).
  37. Bhakta, M. S. et al. A predictive model for time-to-flowering in the common bean based on QTL and environmental variables. *G3-Genes Genom. Genet.* **7**, 3901–3912 (2017).
  38. Marshall-Colon, A. et al. Crops in silico: generating virtual crops using an integrative and multi-scale modeling platform. *Front. Plant Sci.* **8**, 786 (2017).
  39. Zhu, X.-G. et al. Plants in silico: why, why now and what?—an integrative platform for plant systems biology research. *Plant Cell Environ.* **39**, 1049–1057 (2016).
  40. Zhu, X.-G., Wang, Y. U., Ort, D. R. & Long, S. P. *e*-photosynthesis: a comprehensive dynamic mechanistic model of C3 photosynthesis: from light capture to sucrose synthesis. *Plant Cell Environ.* **36**, 1711–1727 (2013).
  41. Kannan, K. et al. Combining gene network, metabolic and leaf-level models shows means to future-proof soybean photosynthesis under rising CO<sub>2</sub>. *in silico. Plants* **1**, diz008 (2019).
  42. Chew, Y. H. et al. Multiscale digital *Arabidopsis* predicts individual organ and whole-organism growth. *Proc. Natl Acad. Sci. USA* **111**, E4127–E4136 (2014).
  43. Xiao, Y. et al. ePlant for quantitative and predictive plant science research in the big data era—Lay the foundation for the future model guided crop breeding, engineering and agronomy. *Quant. Biol.* **5**, 260–271 (2017).
  44. Earles, J. M. et al. Embracing 3D complexity in leaf carbon–water exchange. *Trends Plant Sci.* **24**, 15–24 (2018).
  45. Hansen, J. W. & Jones, J. W. Scaling-up crop models for climate variability applications. *Agr. Syst.* **65**, 43–72 (2000).
  46. Lawrence, D. M. et al. The Community Land Model version 5: description of new features, benchmarking, and impact of forcing uncertainty. *J. Adv. Model. Earth Sy.* **11**, 4245–4287 (2019).
  47. Peng, B. et al. Improving maize growth processes in the community land model: implementation and evaluation. *Agr. Forest Meteorol.* **250–251**, 64–89 (2018).
  48. Scanlon, B. R. et al. The food–energy–water nexus: transforming science for society. *Water Resour. Res.* **53**, 3550–3556 (2017).
  49. Levis, S. et al. Interactive crop management in the Community Earth System Model (CESM1): seasonal influences on land–atmosphere fluxes. *J. Climate* **25**, 4839–4859 (2012).
  50. Osborne, T. et al. JULES-crop: a parametrisation of crops in the Joint UK Land Environment Simulator. *Geosci. Model Dev.* **8**, 1139–1155 (2015).
  51. Wu, X. et al. ORCHIDEE-CROP (v0), a new process-based agro-land surface model: model description and evaluation over Europe. *Geosci. Model Dev.* **9**, 857–873 (2016).
  52. Drewniak, B., Song, J., Prell, J., Kotamarthi, V. R. & Jacob, R. Modeling agriculture in the Community Land Model. *Geosci. Model Dev.* **6**, 495–515 (2013).
  53. Dunbabin, V. M. et al. Modelling root–soil interactions using three-dimensional models of root growth, architecture and function. *Plant Soil* **372**, 93–124 (2013).
  54. Wang, Y. et al. Development of a three-dimensional ray-tracing model of sugarcane canopy photosynthesis and its application in assessing impacts of varied row spacing. *BioEnerg. Res.* **10**, 626–634 (2017).
  55. Vos, J. et al. Functional–structural plant modelling: a new versatile tool in crop science. *J. Exp. Bot.* **61**, 2101–2115 (2009).
  56. Bonan, G. B. et al. Modeling canopy-induced turbulence in the Earth system: a unified parameterization of turbulent exchange within plant canopies and the roughness sublayer (CLM-ml v0). *Geosci. Model Dev.* **11**, 1467–1496 (2018).
  57. Ewert, F. et al. Scale changes and model linking methods for integrated assessment of agri-environmental systems. *Agr. Ecosyst. Environ.* **142**, 6–17 (2011).
  58. Müller, C. et al. Global gridded crop model evaluation: benchmarking, skills, deficiencies and implications. *Geosci. Model Dev.* **10**, 1403–1422 (2017).
  59. Elliott, J. et al. The Global Gridded Crop Model Intercomparison: data and modeling protocols for Phase 1 (v1.0). *Geosci. Model Dev.* **8**, 261–277 (2015).
  60. Rosenzweig, C. et al. The Agricultural Model Intercomparison and Improvement Project (AgMIP): protocols and pilot studies. *Agr. Forest Meteorol.* **170**, 166–182 (2013).
  61. Hoffmann, H. et al. Impact of spatial soil and climate input data aggregation on regional yield simulations. *PLoS ONE* **11**, e0151782 (2016).
  62. Chaney, N. W. et al. Harnessing big data to rethink land heterogeneity in Earth system models. *Hydrol. Earth Syst. Sci.* **22**, 3311–3330 (2018).
  63. Webber, H. et al. Climate change impacts on European crop yields: do we need to consider nitrogen limitation? *Eur. J. Agron.* **71**, 123–134 (2015).
  64. Zimmermann, A. et al. Climate change impacts on crop yields, land use and environment in response to crop sowing dates and thermal time requirements. *Agr. Syst.* **157**, 81–92 (2017).
  65. Boote, K. J., Jones, J. W., White, J. W., Asseng, S. & Lizaso, J. I. Putting mechanisms into crop production models. *Plant Cell Environ.* **36**, 1658–1672 (2013).
  66. Asseng, S. et al. Uncertainty in simulating wheat yields under climate change. *Nat. Clim. Change* **3**, 827–832 (2013).
  67. Bassu, S. et al. How do various maize crop models vary in their responses to climate change factors? *Glob. Change Biol.* **20**, 2301–2320 (2014).
  68. Fleisher, D. H. et al. A potato model intercomparison across varying climates and productivity levels. *Glob. Change Biol.* **23**, 1258–1281 (2017).
  69. Li, T. et al. Uncertainties in predicting rice yield by current crop models under a wide range of climatic conditions. *Glob. Change Biol.* **21**, 1328–1341 (2015).
  70. Rosenzweig, C. et al. Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. *Proc. Natl Acad. Sci. USA* **111**, 3268–3273 (2014).
  71. Martre, P. et al. Multimodel ensembles of wheat growth: many models are better than one. *Glob. Change Biol.* **21**, 911–925 (2015).
  72. Ainsworth, E. A., Leakey, A. D. B., Ort, D. R. & Long, S. P. FACE-ing the facts: inconsistencies and interdependence among field, chamber and modeling studies of elevated [CO<sub>2</sub>] impacts on crop yield and food supply. *New Phytol.* **179**, 5–9 (2008).
  73. Tao, F. et al. Why do crop models diverge substantially in climate impact projections? A comprehensive analysis based on eight barley crop models. *Agr. Forest Meteorol.* **281**, 107851 (2020).
  74. Wang, E. et al. The uncertainty of crop yield projections is reduced by improved temperature response functions. *Nat. Plants* **3**, 17102 (2017).
  75. Wallach, D. et al. Multimodel ensembles improve predictions of crop–environment–management interactions. *Glob. Change Biol.* **24**, 5072–5083 (2018).
  76. Rötter, R. P., Carter, T. R., Olesen, J. E. & Porter, J. R. Crop-climate models need an overhaul. *Nat. Clim. Change* **1**, 175–177 (2011).
  77. Manderscheid, R., Erbs, M. & Weigel, H.-J. Interactive effects of free-air CO<sub>2</sub> enrichment and drought stress on maize growth. *Eur. J. Agr.* **52**, 11–21 (2014).
  78. Ainsworth, E. A. & Long, S. P. What have we learned from 15 years of free-air CO<sub>2</sub> enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO<sub>2</sub>. *New Phytol.* **165**, 351–372 (2005).
  79. Kimball, B. A. Lessons from FACE: CO<sub>2</sub> effects and interactions with water, nitrogen, and temperature. *Curr. Opin. Plant Biol.* **31**, 36–43 (2010).
  80. Kimball, B. A. Crop responses to elevated CO<sub>2</sub> and interactions with H<sub>2</sub>O, N, and temperature. *Curr. Opin. Plant Biol.* **31**, 36–43 (2016).
  81. Bernacchi, C. J., Kimball, B. A., Quarles, D. R., Long, S. P. & Ort, D. R. Decreases in stomatal conductance of soybean under open-air elevation of [CO<sub>2</sub>] are closely coupled with decreases in ecosystem evapotranspiration. *Plant Physiol.* **143**, 134–144 (2007).
  82. Gray, S. B. et al. Intensifying drought eliminates the expected benefits of elevated carbon dioxide for soybean. *Nat. Plants* **2**, 16132 (2016).
  83. Jin, Z., Ainsworth, E. A., Leakey, A. D. B. & Lobell, D. B. Increasing drought and diminishing benefits of elevated carbon dioxide for soybean yields across the US Midwest. *Glob. Change Biol.* **24**, e522–e533 (2018).
  84. Sanz-Sáez, Á. et al. Leaf and canopy scale drivers of genotypic variation in soybean response to elevated carbon dioxide concentration. *Glob. Change Biol.* **23**, 3908–3920 (2017).
  85. Bishop, K. A., Betzelberger, A. M., Long, S. P. & Ainsworth, E. A. Is there potential to adapt soybean (*Glycine max* Merr.) to future [CO<sub>2</sub>]? An analysis of the yield response of 18 genotypes in free-air CO<sub>2</sub> enrichment. *Plant Cell Environ.* **38**, 1765–1774 (2015).
  86. Ainsworth, E. A. & Rogers, A. The response of photosynthesis and stomatal conductance to rising [CO<sub>2</sub>]: mechanisms and environmental interactions. *Plant Cell Environ.* **30**, 258–270 (2007).

87. Cai, C. et al. Responses of wheat and rice to factorial combinations of ambient and elevated CO<sub>2</sub> and temperature in FACE experiments. *Glob. Change Biol.* **22**, 856–874 (2016).
88. Ruiz-Vera, U. M., Siebers, M. H., Drag, D. W., Ort, D. R. & Bernacchi, C. J. Canopy warming caused photosynthetic acclimation and reduced seed yield in maize grown at ambient and elevated [CO<sub>2</sub>]. *Glob. Change Biol.* **21**, 4237–4249 (2015).
89. Sinclair, T. R. & Muchow, R. C. in *Advances in Agronomy* Vol. 65 (Ed. Sparks, D. L.) 215–265 (Academic Press, 1999).
90. Yin, X. & Struiik, P. C. Can increased leaf photosynthesis be converted into higher crop mass production? A simulation study for rice using the crop model GECROS. *J. Exp. Bot.* **68**, 2345–2360 (2017).
91. Vanuytrecht, E. & Thorburn, P. J. Responses to atmospheric CO<sub>2</sub> concentrations in crop simulation models: a review of current simple and semicomplex representations and options for model development. *Glob. Change Biol.* **23**, 1806–1820 (2017).
92. Huntingford, C. et al. Implications of improved representations of plant respiration in a changing climate. *Nat. Commun.* **8**, 1602 (2017).
93. Yin, X. Improving ecophysiological simulation models to predict the impact of elevated atmospheric CO<sub>2</sub> concentration on crop productivity. *Annal. Bot.* **112**, 465–475 (2013).
94. Asseng, S., Kassie, B. T., Labra, M. H., Amador, C. & Calderini, D. F. Simulating the impact of source-sink manipulations in wheat. *Field Crop. Res.* **202**, 47–56 (2017).
95. Emberson, L. D. et al. Ozone effects on crops and consideration in crop models. *Eur. J. Agr.* **100**, 19–34 (2018).
96. Ewert, F. & Porter, J. R. Ozone effects on wheat in relation to CO<sub>2</sub>: modelling short-term and long-term responses of leaf photosynthesis and leaf duration. *Glob. Change Biol.* **6**, 735–750 (2000).
97. Guarin, J. R., Kassie, B., Mashaeet, A. M., Burkey, K. & Asseng, S. Modeling the effects of tropospheric ozone on wheat growth and yield. *Eur. J. Agr.* **105**, 13–23 (2019).
98. van Oijen, M., Dreccer, M. F., Firsching, K. H. & Schnieders, B. J. Simple equations for dynamic models of the effects of CO<sub>2</sub> and O<sub>3</sub> on light-use efficiency and growth of crops. *Ecol. Model.* **179**, 39–60 (2004).
99. Ainsworth, E. A., Yendrek, C. R., Sitch, S., Collins, W. J. & Emberson, L. D. The effects of tropospheric ozone on net primary productivity and implications for climate change. *Annual Rev. Plant Biol.* **63**, 637–661 (2012).
100. Tao, F., Feng, Z., Tang, H., Chen, Y. & Kobayashi, K. Effects of climate change, CO<sub>2</sub> and O<sub>3</sub> on wheat productivity in Eastern China, singly and in combination. *Atmos. Environ.* **153**, 182–193 (2017).
101. Field, C. B., Barros, V., Stocker, T. F. & Dahe, Q. Managing the risks of extreme events and disasters to advance climate change adaptation: special report of the intergovernmental panel on climate change. (Cambridge Univ. Press, 2012).
102. Lesk, C., Rowhani, P. & Ramankutty, N. Influence of extreme weather disasters on global crop production. *Nature* **529**, 84–87 (2016).
103. Barnabás, B., Jäger, K. & Fehér, A. The effect of drought and heat stress on reproductive processes in cereals. *Plant Cell Environ.* **31**, 11–38 (2008).
104. Eyshi Rezaei, E., Webber, H., Gaiser, T., Naab, J. & Ewert, F. Heat stress in cereals: mechanisms and modelling. *Eur. J. Agr.* **64**, 98–113 (2015).
105. Prasad, P. V. V., Bheemanahalli, R. & Jagadish, S. V. K. Field crops and the fear of heat stress—Opportunities, challenges and future directions. *Field Crop. Res.* **200**, 114–121 (2017).
106. Shi, W. et al. High day- and night-time temperatures affect grain growth dynamics in contrasting rice genotypes. *J. Exp. Bot.* **68**, 5233–5245 (2017).
107. Peng, S. et al. Rice yields decline with higher night temperature from global warming. *Proc. Natl Acad. Sci. USA* **101**, 9971–9975 (2004).
108. Saini, H. S. & Westgate, M. E. in *Advances in Agronomy* Vol. 68 (Ed. Sparks, D. L.) 59–96 (Academic Press, 1999).
109. Mazdiyasi, O. & AghaKouchak, A. Substantial increase in concurrent droughts and heatwaves in the United States. *Proc. Natl Acad. Sci. USA* **112**, 11484–11489 (2015).
110. Lobell, D. B. et al. The shifting influence of drought and heat stress for crops in northeast Australia. *Glob. Change Biol.* **21**, 4115–4127 (2015).
111. Liu, B. et al. Testing the responses of four wheat crop models to heat stress at anthesis and grain filling. *Glob. Change Biol.* **22**, 1890–1903 (2016).
112. Barlow, K. M., Christy, B. P., O’Leary, G. J., Riffkin, P. A. & Nuttall, J. G. Simulating the impact of extreme heat and frost events on wheat crop production: a review. *Field Crop. Res.* **171**, 109–119 (2015).
113. Siebert, S., Webber, H., Zhao, G. & Ewert, F. Heat stress is overestimated in climate impact studies for irrigated agriculture. *Environ. Res. Lett.* **12**, 054023 (2017).
114. Webber, H. et al. Diverging importance of drought stress for maize and winter wheat in Europe. *Nat. Commun.* **9**, 4249 (2018).
115. Siebert, S., Ewert, F., Rezaei, E. E., Kage, H. & Graß, R. Impact of heat stress on crop yield—the importance of considering canopy temperature. *Environ. Res. Lett.* **9**, 044012 (2014).
116. Webber, H. et al. Physical robustness of canopy temperature models for crop heat stress simulation across environments and production conditions. *Field Crop. Res.* **216**, 75–88 (2018).
117. Rosenzweig, C., Tubiello, F. N., Goldberg, R., Mills, E. & Bloomfield, J. Increased crop damage in the US from excess precipitation under climate change. *Glob. Environ. Change* **12**, 197–202 (2002).
118. Ebrahimi-Mollabashi, E. et al. Enhancing APSIM to simulate excessive moisture effects on root growth. *Field Crop. Res.* **236**, 58–67 (2019).
119. Li, Y., Guan, K., Schnitkey, G. D., DeLucia, E. & Peng, B. Excessive rainfall leads to maize yield loss of a comparable magnitude to extreme drought in the United States. *Glob. Change Biol.* **25**, 2325–2337 (2019).
120. Constantin, J. et al. Management and spatial resolution effects on yield and water balance at regional scale in crop models. *Agr. Forest Meteorol.* **275**, 184–195 (2019).
121. Brilli, L. et al. Review and analysis of strengths and weaknesses of agro-ecosystem models for simulating C and N fluxes. *Sci. Total Environ.* **598**, 445–470 (2017).
122. Luo, Y. et al. Toward more realistic projections of soil carbon dynamics by Earth system models. *Global Biogeochem. Cy.* **30**, 40–56 (2015).
123. Koven, C. D. et al. The effect of vertically resolved soil biogeochemistry and alternate soil C and N models on C dynamics of CLM4. *Biogeochemistry* **10**, 7109–7131 (2013).
124. Tang, J. Y., Riley, W. J., Koven, C. D. & Subin, Z. M. CLM4-BeTR, a generic biogeochemical transport and reaction module for CLM4: model development, evaluation, and application. *Geosci. Model Dev.* **6**, 127–140 (2013).
125. Niu, S. et al. Global patterns and substrate-based mechanisms of the terrestrial nitrogen cycle. *Ecol. Lett.* **19**, 697–709 (2016).
126. Rötter, R. P. et al. Simulation of spring barley yield in different climatic zones of Northern and Central Europe: a comparison of nine crop models. *Field Crop. Res.* **133**, 23–36 (2012).
127. Palosuo, T. et al. Simulation of winter wheat yield and its variability in different climates of Europe: a comparison of eight crop growth models. *Eur. J. Agr.* **35**, 103–114 (2011).
128. Ehrhardt, F. et al. Assessing uncertainties in crop and pasture ensemble model simulations of productivity and N<sub>2</sub>O emissions. *Glob. Change Biol.* **24**, e603–e616 (2018).
129. Basso, B. et al. Soil organic carbon and nitrogen feedbacks on crop yields under climate change. *Agricultural & Environmental Letters* **3**, 180026 (2018).
130. Basso, B., Hyndman, D. W., Kendall, A. D., Grace, P. R. & Robertson, G. P. Can impacts of climate change and agricultural adaptation strategies be accurately quantified if crop models are annually re-initialized? *PLoS ONE* **10**, e0127333 (2015).
131. Kollas, C. et al. Crop rotation modelling—a European model intercomparison. *Eur. J. Agr.* **70**, 98–111 (2015).
132. McDermid, S., Mearns, L. & Ruane, A. Representing agriculture in Earth system models: approaches and priorities for development. *J. Adv. Model. Earth Sy.* **9**, 2230–2265 (2017).
133. Deutsch, C. A. et al. Increase in crop losses to insect pests in a warming climate. *Science* **361**, 916–919 (2018).
134. Savary, S. et al. Crop health and its global impacts on the components of food security. *Food Secur.* **9**, 311–327 (2017).
135. Porter, J. H., Parry, M. L. & Carter, T. R. The potential effects of climatic change on agricultural insect pests. *Agr. Forest Meteorol.* **57**, 221–240 (1991).
136. Donatelli, M. et al. Modelling the impacts of pests and diseases on agricultural systems. *Agr. Syst.* **155**, 213–224 (2017).
137. Lammoglia, S.-K. et al. Modelling pesticides leaching in cropping systems: effect of uncertainties in climate, agricultural practices, soil and pesticide properties. *Environ. Modell. Softw.* **109**, 342–352 (2018).
138. Wang, R. et al. A review of pesticide fate and transport simulation at watershed level using SWAT: current status and research concerns. *Sci. Total Environ.* **669**, 512–526 (2019).
139. Ruane, A. C. et al. An AgMIP framework for improved agricultural representation in IAMs. *Environ. Res. Lett.* **12**, 125003 (2017).
140. Rötter, R. P. et al. Linking modelling and experimentation to better capture crop impacts of agroclimatic extremes—a review. *Field Crop. Res.* **221**, 142–156 (2018).
141. Schlenker, W. & Roberts, M. J. Nonlinear temperature effects indicate severe damages to U. S. crop yields under climate change. *Proc. Natl Acad. Sci. USA* **106**, 15594–15598 (2009).
142. Grunwald, S., Thompson, J. & Boettinger, J. Digital soil mapping and modeling at continental scales: finding solutions for global issues. *Soil Sci. Soc. Am. J.* **75**, 1201–1213 (2011).
143. Hengl, T. et al. SoilGrids1km—global soil information based on automated mapping. *PLoS ONE* **9**, e105992 (2014).

144. Chaney, N. W. et al. POLARIS soil properties: 30-meter probabilistic maps of soil properties over the contiguous United States. *Water Resour. Res.* **55**, 2916–2938 (2019).
145. Han, E., Ines, A. V. M. & Koo, J. Development of a 10-km resolution global soil profile dataset for crop modeling applications. *Environ. Modell. Softw.* **119**, 70–83 (2019).
146. Coucheney, E. et al. Key functional soil types explain data aggregation effects on simulated yield, soil carbon, drainage and nitrogen leaching at a regional scale. *Geoderma* **318**, 167–181 (2018).
147. Pongratz, J. et al. Models meet data: challenges and opportunities in implementing land management in Earth system models. *Glob. Change Biol.* **24**, 1470–1487 (2018).
148. Gbегbelegbe, S. et al. Baseline simulation for global wheat production with CIMMYT mega-environment specific cultivars. *Field Crop. Res.* **202**, 122–135 (2017).
149. Woodard, J. D. et al. The power of agricultural data. *Science* **362**, 410–411 (2018).
150. Minet, J. et al. Crowdsourcing for agricultural applications: a review of uses and opportunities for a farmsourcing approach. *Comput. Electron. Agr.* **142**, 126–138 (2017).
151. van Bussel, L. G. J., Ewert, F. & Leffelaar, P. A. Effects of data aggregation on simulations of crop phenology. *Agr. Ecosyst. Environ.* **142**, 75–84 (2011).
152. Boryan, C., Yang, Z., Mueller, R. & Craig, M. Monitoring US agriculture: the US department of agriculture, national agricultural statistics service, cropland data layer program. *Geocarto Int.* **26**, 341–358 (2011).
153. Xie, Y., Lark, T. J., Brown, J. F. & Gibbs, H. K. Mapping irrigated cropland extent across the conterminous United States at 30 m resolution using a semi-automatic training approach on Google Earth Engine. *ISPRS J. Photogramm.* **155**, 136–149 (2019).
154. Azzari, G. et al. Satellite mapping of tillage practices in the North Central US region from 2005 to 2016. *Remote Sens. Environ.* **221**, 417–429 (2019).
155. Seifert, C. A., Azzari, G. & Lobell, D. B. Satellite detection of cover crops and their effects on crop yield in the Midwestern United States. *Environ. Res. Lett.* **13**, 064033 (2018).
156. Urban, D., Guan, K. & Jain, M. Estimating sowing dates from satellite data over the U. S. Midwest: a comparison of multiple sensors and metrics. *Remote Sens. Environ.* **211**, 400–412 (2018).
157. Lobell, D. B., Sibley, A. & Ortiz-Monasterio, J. I. Extreme heat effects on wheat senescence in India. *Nat. Clim. Change* **2**, 186–189 (2012).
158. Sakamoto, T. et al. A two-step filtering approach for detecting maize and soybean phenology with time-series MODIS data. *Remote Sens. Environ.* **114**, 2146–2159 (2010).
159. Baldocchi, D. et al. FLUXNET: A new tool to study the temporal and spatial variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities. *B. Am. Meteorol. Soc.* **82**, 2415–2434 (2001).
160. Kimball, B. A. et al. Simulation of maize evapotranspiration: an inter-comparison among 29 maize models. *Agr. Forest Meteorol.* **271**, 264–284 (2019).
161. Boote, K. J., Prasad, V., Allen, L. H. Jr, Singh, P. & Jones, J. W. Modeling sensitivity of grain yield to elevated temperature in the DSSAT crop models for peanut, soybean, dry bean, chickpea, sorghum, and millet. *Eur. J. Agr.* **100**, 99–109 (2017).
162. Lobell, D. B. et al. Greater sensitivity to drought accompanies maize yield increase in the U. S. Midwest. *Science* **344**, 516–519 (2014).
163. Lobell, D. B. & Asseng, S. Comparing estimates of climate change impacts from process-based and statistical crop models. *Environ. Res. Lett.* **12**, 015001 (2017).
164. Roberts, M. J., Braun, N. O., Sinclair, T. R., Lobell, D. B. & Schlenker, W. Comparing and combining process-based crop models and statistical models with some implications for climate change. *Environ. Res. Lett.* **12**, 095010 (2017).
165. Guan, K. et al. The shared and unique values of optical, fluorescence, thermal and microwave satellite data for estimating large-scale crop yields. *Remote Sens. Environ.* **199**, 333–349 (2017).
166. Luo, Y., Guan, K. & Peng, J. STAIR: a generic and fully-automated method to fuse multiple sources of optical satellite data to generate a high-resolution, daily and cloud-/gap-free surface reflectance product. *Remote Sens. Environ.* **214**, 87–99 (2018).
167. Viña, A., Gitelson, A. A., Nguy-Robertson, A. L. & Peng, Y. Comparison of different vegetation indices for the remote assessment of green leaf area index of crops. *Remote Sens. Environ.* **115**, 3468–3478 (2011).
168. Anderson, M. et al. Mapping daily evapotranspiration at field to continental scales using geostationary and polar orbiting satellite imagery. *Hydrol. Earth Syst. Sc.* **15**, 223–239 (2011).
169. Cai, Y. et al. Integrating satellite and climate data to predict wheat yield in Australia using machine learning approaches. *Agr. Forest Meteorol.* **274**, 144–159 (2019).
170. Huang, J. et al. Assimilation of remote sensing into crop growth models: current status and perspectives. *Agr. Forest Meteorol.* **276–277**, 107609 (2019).
171. Asseng, S. et al. Model-driven multidisciplinary global research to meet future needs: the case for “improving radiation use efficiency to increase yield”. *Crop Sci.* **59**, 843–849 (2019).
172. Vermeulen, S. et al. Climate change, agriculture and food security: a global partnership to link research and action for low-income agricultural producers and consumers. *Curr. Opin. Env. Sust.* **4**, 128–133 (2012).

### Acknowledgements

B.P., K.G., H.K. and W.Z. are supported by the United States National Science Foundation (NSF) Career Award (grant no. 1847334), National Aeronautics and Space Administration (NASA) Carbon Monitoring System managed by NASA Terrestrial Ecology Program (grant no. 80NSSC18K0170), United States Department of Agriculture (USDA) National Institute of Food and Agriculture (NIFA) Program (grant no. 2017-67013-26253) to K.G. B.P., K.G., S.A. and D.I.G. acknowledge support by USDA NIFA (grant no. 2017-68002-26789). B.P., K.G., W.Z., A.M.C. and J.C.S. acknowledge support by the Foundation for Food and Agriculture Research (FFAR) (grant no. 602757). B.P., K.G., D.M.L. and D.L.L. acknowledge support by the National Center for Atmospheric Research, which is a major facility sponsored by the NSF under cooperative agreement no. 1852977. The content of this publication is solely the responsibility of the authors and does not necessarily represent the official views of the FFAR. Any opinions, findings and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the USDA. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the USDA. Due to space constraints, we could not cite all relevant literature. We apologize to the authors whose important work was not cited in this Perspective.

### Author contributions

B.P. and K.G. conceived the research. B.P., K.G. and J.T. wrote the paper; B.P. and K.G. designed the figures. B.P. and H.K. produced the figures. B.P. designed and produced the table. E.A.A., S.A., C.J.B., M.C., E.H.D., J.W.E., F.E., R.F.G., D.I.G., G.L.H., J.W.J., Z.J., H.K., D.M.L., Y.L., D.L.L., A.M.C., C.D.M., D.R.O., J.C.S., C.E.V., A.W., X.Y. and W.Z. all contributed to the text.

### Competing interests

The authors declare no competing interests.

### Additional information

Correspondence should be addressed to B.P. or K.G.

Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© Springer Nature Limited 2020